INDEX

Acer negundo, 422, 423, 424.
platanoides, 422, 423, 424.
rubrum, 421, 422, 423, 424.
saccharinum, 421.
saccharum, 421, 423, 424.
spicatum, 422.
Achras sapotes, 156.
Acrobasis caryæ, 197, 199.
Acrobatic fly, 330.
Adalia bipunctata, 404, 405, 408.
Agallia uhleri, 426.
Agropyron biflorum, 143.
Alabama argillacea, in Peru, 244–46.
clover, 362.
Alfalfa, 189, 192, 194.
butterfly, 192.
gall midge, 194.
Allorhina mutabilis, 193.
nitida, 193.
American Association Economic Entomologists, VII–XVI.
elm, 285.
Anasa tristis, 190.
Anastrepha fratercula, 345.
peruviana, 345.
Anisocalvia duodecimmaculata, 406.
quatuordecimguttata, 406.
Anthocoptes transitionalis, 420.
Anthonomus vestitus, 10, 11, 303, 304.
Ants, 195.
Aphelinus fuscipennis, 144, 273, 320.
mytilaspidis, 144.
quaylei, 320.
Aphis, 10.
avenæ, 406.
brassicae, 190.
gossypii, 190.
pomi, 406.
sorbi, 406, 457.
viburni, 406.
Aphycus flavus, 301.
sp., 294, 295, 300.
Apiculture, means of promoting, 418–20.
Apple, 62, 64, 65, 66, 76, 157, 158, 160, 166, 167, 168, 169, 170, 174, 175, 176, 177, 178, 193, 240, 313, 316, 390, 396, 412.
Apple and cherry ermine moths, 63.
leaf-hopper, 240–43.
maggots, 79.
scab, 165, 177.
Apricot, 263.
Archips argyrospila, pupal instar of, 396–98.
Archytas piliventris, 235, 257.
Argas persicus, 96.
Argentine ant, 294.
Arrhenophagus chionaspidis, 322.
Arsenate of lead, 61, 80, 157, 158, 159, 166, 365, 478.
powdered, 477–79.
Arsenic, white, 440.
Arsenical residues on fruit, 159.
Arsenite of zinc, 365.
Arthrocnodax carolina, 488–89.
occidentalis, 87.
Artichoke, 178.
Ash, 76, 78, 178.
Asphondylia capparis, 279.
miki, 194.
Aspidiotiphagus citrinus, 319.
Aspidiotus ostreæformis, 144.
rapax, 144.
Aster, 180, 238, 410.
Australian lady-bird beetle, 256.
Autographa brassicæ, 144, 145.

Bacillus amylovorus, 178.
Bananas, 68, 70, 71, 73, 328.
Barbiero fever, 105.
Barley, 117.
Bean, 189, 190, 325.
lady bug, 193.
Bed-bug, 104, 109, 189, 217, 382.
Bees, 187, 418.
Beech, 410.
Bell pepper, 362.
Bermuda grass, 362.
Birch, 410.
Birds, 255, 360.
Bird lice, 82.
Bishopp, F. C., 112–26.
and King, W. V., 200–11.
Bitter-rot, 166, 170, 173.
Blackberry, 178, 410.
Blackbirds, 236, 254, 257.
Black cherry aphis, 406, 457.
leaf 40, 65, 414.
locust, 178.
fly, 112.
maple, 422, 423, 424.
scale, 293, 294, 300, 301.
wasp, 456.
Bleeding trees, 285.
Blight canker, 178.
Blood sucking cone nose, 189.
Blow fly, 111.
Blue grass, 362.
Boll weevil, 57.
Bordeaux, 157, 158, 166, 173, 454.
Bot flies, 111.

Box elder, 178.
Bran, 236.
Britton, W. E., 89–93; 197–99.
Brown lace wing, 87.
Brown-tail moth, 74, 79, 134, 136, 159, 331, 332; attempts to trap, 74.
Bruchophagus funebris, 194.
Brues, C. T., 101–109, 459–77.
Bud moth, 198, 199.
weevils, 12.
Buffalo gnats, 217.
Burgess, A. F., 258–59; 286, 427.
and Rogers, D. M., 75–79.
Butternut, 177.
Buttrick, P. L., 352–59.

Cabbage, 157.
louse, 190.
Cacao canker, 442.
pods, 441, 442.
California red scale, 191, 192.
Calosoma alternans, 235.
calidum, 364.
Camphor tree, 413, 414.
Campylomma verbasci, 64.
Cantharoctonus stramineus, 308.
Capparis ægyptiaca, 279.
spinosa, 279.
Campsomeris dorsata, 254, 255.
Carabid beetle, 235.
Cardin, Patricio, 330.
Carolina poplar, 412.
Carpocapsa pomonella, 192.
Carpophilus dimidiatus, 193.
Cassaba melons, 190.
Cassava starch, 452.
Castnia licus, 448, 453.
Castolus plagiaticollis, 247, 453.
Castor bean, 325.
Catalpa, 178.
Catolaccus townsendi, 308, 309.
Cecidomyia peritomatis, 280.
Cerambycobius peruvianus, 308.
townsendi, 308.
Ceratitis capitata, 68, 70, 347.
Ceratophyllus gallinæ, 84.
Ceratopogon hippocastani, 285.
Cerchysius sp., 301.
Ceresa bubalus, 412.
taurina, 411, 412.
Changa, 149.
Charcoal, 456.
Chelonella townsendi, 308.
Cherry, 62, 64, 66, 79, 80, 81, 177, 178, 306.
fruit-flies, 79–80.
sawfly leaf-miner, 62.
Chestnut, 76, 77, 78, 177.
bark disease, 281, 404, 405.
Chilocorus bivulnerus, 273.
similis, 224, 322.
China tree, 191.
Chinch bug parasite, 425, 489.
Chinese banana, 400.
Cholera, 32, 35.
Chrysomphalus aurantii, 191, 299.
Cigarette beetle, 226.
Cirphis latiuscula, 257.
Citellus beecheyi, 205.
columbianus, 209.
Citrus, 85, 86, 88, 150, 161, 189, 191, 192, 293, 345, 347.
buxifolius, 401.
medica limonum, 400.
red spider, 87.
thrips, 186, 189.
Clover, 238, 278, 283.
leaf weevil, 284.
seed chalcis, 194.
Coccinella novemnotata, 404, 406, 408.
transversoguttata, 404, 406.
Coccophagus lecanii, 294, 295, 298, 299, 300, 303.
lunulatus, 294, 295, 299, 300.
Coccus hesperidum, 11; parasites of, 293–303.
sacchari, 252.
Cockerell, T. D. A., 142–43; 279–80; 424, 425, 426, 489.
Cockle bur, 362.
Cockroach trap, 327–29.
Coconuts, 442.
Codling moth, 36, 166, 170, 171, 173, 175, 176, 192, 270, 389; brood study of, 389–95; one spray, 173–75.
Coffee, 150.
ant, 150.
Coloradо potato beetle, 193.
Compsilura concinnata, 485.
Confused flour beetle, 40.
Conorhinus sanguisugus, 189.
Conotrachelus juglandis, 197.
Conventzia hageni, 85.
Cooley, R. A., 93–95.
Cordyceps barberi, 254.
Corn, 30, 231, 278, 362, 365, 425, 426, 445.
ear worm, 192.
silk, 342.
Corthylus punctatissimus, 427.
Corythuca ciliata, 407–8.
Cosmocomoidea morrilli, 249.
Cotton, 11, 36, 245, 304, 305, 307, 309, 310, 312, 318, 319, 325, 326, 362, 365.
boll weevil, 311.
insects, 11.
leaf-caterpillar, 244.
moth flight, 426–7.
square-weevil, 10, 303.
worm, 478.
Cottonwood, 178.
Cow pea, 362, 365.
weevil, 147.
manure, 463.
Crab apple, 412.
grass, 362.
Cricket, 150.

Crotophaga ani, 236, 255.
Crown gall, 267.
Crow, 365.
Cryptolæmus montrouzieri, 12, 256.
Cryptothrips floridensis, 413–14.
Culex pipiens, 93.
sollicitans, 91, 352.
Curculio, 166, 169.
Currant, 178.
Cut-worm moths, 267.
Cyanopepla submacula, 249.
Cycloneda sanguinea, 149, 257, 404.
Cypress, 278.

Dacus zonatus, 348.
Daictimorpha peruviana, 308.
Disprepes abbreviatus, 455.
Dahlia, 410.
Daisy, 238.
Date palm, 192, 193, 415.
scales, 186, 192.
Daucus carota, 180.
Davidson, W. M., 396–98.
Davis, J. J., 276–78.
Dean, G. A., 40–53.
Deer fly, 112.
Delphax saccharivora, 251, 252, 256.
Demerara, Trinidad and Barbadoes trip, 443–57.
Dengue, 31, 104.
Dermacentor venustus, 93, 200, 210.
vulpinus, 55.
Dew, J. A., 361–66.
Diabrotica graminea, 257.
Diaprepes abbreviatus, 256.
Diarrhea, 34, 107.
Diaspis pentagona, 321, 322.
Diatræa canella, 445.
grandiosella, 247, 248.
lineolata, 445.
saccharalis, 12, 247, 251, 252, 253, 254, 445.
Dicraneura cockerelli, 191.
Distillate oil, 66.
Doane, R. W., 366–85, 437–42.
Dogwood, 178, 410.
Dolichoderus bituberculatus, 143.
Doryphora decimlineata, 193.
Douglas fir-seed chalcid, 283.
Drosicha contrahens, 143.
corpulenta, 143.
lichenoides, 142, 143.
maskelli, 143.
stebbingi, 143.
Dyar, H. G., 336.
Dysentery, 32, 34.

Eastern grape leaf skeletonizer, 192.
peach tree borer, 193, 263, 264.
rose chafer, 193.
Econonomic Entomologists, 26th annual meeting, 427.
Proceedings, 1–126; 157–257; 293–329.
Economic Entomology, Second International Congress, 88.
Ectatomma tuberculatum, 309, 312.
Egyptian cotton, 190.
Eight-lined Gypona, 409.
Elder, 178.
Elis sexcincta, 254, 255.
Elm, 157, 177, 178.
Emposasca mali, 240.
English sparrow, 365.
walnut, 197.
Eniscopilus purgatus, 365.
Enteritis, 33, 34.
Entomological bulletin, arrangement of material, 180–84.
conferences in Porto Rico, 148–50.
nomenclature, 340.
pioneering, 185–95.
Society of Ontario, 150.
Ephestia kuehniella, 144.
Epilachna corrupta, 193.
Ericoccus borealis, 424.
Erigeron canadensis, 178, 238.
Eriophyes confusus, 421.
elongatus, 421.
maculatus, 421.
major, 421.
modestus, 421.
negundi, 422.
ornatus, 422.
parallelus, 422.
regulus, 422.
Ermine moth, 66.
Escherich, K., 430.
Essig, E. O., 270–73; 289.
Eucelatoria australis, 246.
European grain aphis, 406.
Eurytoma piuræ, 308.
Eusemion longipenne, 294, 301, 302.
Eutettix tenella, 191.
Euthrips citri, 186.
pyri, 62.
tritici, 187.
Euvanessa antiopa, 157.
Ewing, H. E., 404–7.

Fall army worm, 230, 361–66.
webworm, 157.
False tarnished plant-bug, 64.
Federal Horticultural Board, 340.
plant quarantine act, 133–42.
Felt, E. P., 160–61; 173–75; 278–79; 283, 284, 331, 427, 488–89.
Ficus nota, 143.
Field larks, 365.
Fig, 191, 193.
eater, 193.
Filaria, 219.
Filariasis, 27, 371.
Fire-ant, 256.
Fish oil, 126.
soap, 244.
Flax, 470.
Flea, 81, 104, 109, 217, 378.

Flour, 43, 44.
Flower or grain thrips, 189.
Folsom, J. W., 429.
Forbes, R. H., 415–16.
Forest entomology, society, 282.
Foul brood, 187.
Fowl bug, 81.
Fowls, domestic, external insect parasites of, 81–84.
Froghopper, 444, 451, 452, 453, 454, 455.
Frontina archippivora, 235, 257.
Fruit tree leaf roller, 396.

Gadflies, 105.
Gasoline torch treatment of date palm scales, 415–16.
Gates, B. E., 418–20.
Geotropism, negative, 407–9.
Giant moth borer, 448, 449, 451.
Gipsy moth, 65, 75, 76, 78, 134, 136, 141, 298, 477; conference, 286; experiments in controlling, 75–79; remarks, 258–59; work, reorganization of, 331; wilt disease, 479–88.
Glaser, R. W., and Chapman, J. W., 479–88.
Glossina morsitans, 367.
palpalis, 367.
Glucose, 157.
Goldenrod, 178, 180.
Gonia crassicornis, 235, 257.
Gouty pine midge, 331.
Grain, 55, 463.
Grape, 64, 178.
leaf hopper, 191, 240.
phylloxera, 190.
Grasshoppers, 248.
mice, 224.
Grass looper, 236.
Grass worms, 231, 232, 234, 236.
Gray birch, 76, 78.
Green apple aphis, 406.
June beetle, 193.
muscardine disease, 453.
peppers, 402.
Ground squirrel, 209.
Gru-gru worm, 454.
Gryllodes muticus, 150.
Gryllotalpa hexadactyla, 252, 253.
Guavas, 156, 345, 346.
Gypona cana, 412.
flavilineata, 410.
octolineata, 409–13.
Gyrococcus flaccidiflez, 480.

Hadley, C. H., Jr., 74.
Hæmatobia, 105.
serrata, 124.
Harrisina americana, 192.
coracina, 192.
metallica, 192.
Harvester ant, 194.
Haseman, Leonard, 237–40; 240–43; 313–16.
Harte, C. R., 426–7.
Hawthorn, 64, 177.
Headlee, T. J., 130–33; 389–95.
Heart-water, 110.
Heat as means of controlling mill insects, 40–53.
Heidemann, Otto, 249–51.
Heliothis armigera, 192.
Hemerobius californicus, 87.
Hemichionaspis aspidistræ, 322.
minor, 9, 11, 318, 319, 320, 321, 322.
Hemisarcoptes malus, 320.
Hemlock, 78.
Hen flea, 84.
louse, 82.
Herms, W. B., 430.
Herrick, G. W., 79–80; 81–84; 152.
Hessian fly, 36.
Heterocordylus malinus, 64.
Heterogeomys hispidus, 248.
Hewitt, C. G., 274–76.
Hickory, 76.
Hinds, W. E., 477–79.
Hippodamia convergens, 404, 405, 406.
parenthesis, 404.
spuria, 404, 406.
Hodge, C. F., 110–12.
Hodge fly trap, 328, 329.
Hodgkiss, H. E., 243–44; 420–24.
Holloway, T. E., 341–44.
Holoquiscalus brachipterus, 236, 255.
Home-made kerosene emulsion, 242.
Homœsoma electellum, 426.
Honey locust, 178.
Hooker, C. W., 148–50; (Obituary), 334; (Writings), 336.
Horsechestnut, 285.
Horismenus urichi, 249.
Horn-fly, 105, 111, 124.
Horse flies, 217.
manure, 117, 124.
weed, 178.
Horticultural inspectors, proceedings, 127–42; 258–69.
House fly, 28, 34, 35, 105, 109, 111, 124, 193, 195, 367, 372, 460.
How contact insecticides kill, 160.
Howard, L. O., 88, 336.
Hungerford, H. B., 165–73.
Hunter, S. J., 96–99; 196–97.
Hunter, W. D., 27–39.
Hydrocyanic acid gas, 40, 41, 44, 45, 53, 54.
Hypera punctata, 284.
Hyphantria cunea, 158.

Ihle, Paul, 430.
Imagination and facts, 147–48.
Imperial Bureau of Entomology, 274–76.
Indian cattle, 32.
Insect collections, Kansas University, 283.

Insect and disease, literature, 366–85.
bibliography, 368–85.
Inspection, What besides, 265–69.
Iridomyrmex humilis, 294.
Irish potatoes, 188, 193, 362.
Iron sulphide, 159.
Italian pear scale, 164.
Itonida anthici, 278–79.
inopis, 331.
Ixodes æqualis, 205.

Johnson grass, 362.
Johnston, F. A., 144–47.
Jonathan tree, 411.
Jones, T. H., 230–36.
Juglans regia, 197.
Juniper, 76.

Kaffir corn, 362.
Kala-azar fever, 104.
Kerosene, 90, 91, 347, 348, 349, 350, 440.
Kerosene emulsion, 65, 163, 164.
Knab, F., 336.

Lachnosterna arcuata, 276.
grubs, 455.
tristis, life cycle of, 276–78.
Lady-bird beetle, 12, 149, 257.
Laphygma frugiperda, 252, 257, 362; in Porto Rico, 230–36.
Larger moth borer, 455.
Lasioderma serricorne, 226.
Lead arsenate, 166, 176, 198; paste, 157.
Leaf cutter bees, 194, 195, 425.
miner, 150.
Leather beetle, 55.
Lecanium corni, 295, 298, 299.
hesperidum, 186, 191.
Leishmania, 219.
Lemon, 399, 400.
Leprosy, 32, 371.
Leptodictya tabida, 248, 249.
Leptoglossus zonatus, 186, 190.
Lesser apple worm, 176.
Leucoptera coffeella, 150.
Lice, 104, 109, 217, 384.
Ligyrus rugiceps, 255.
Lilac, 177.
Lime-sulphur, 157, 158, 164, 173, 244, 266, 270, 271, 272.
Lina scripta, 158.
Limotettix exitiosus, 426.
Liquid chocolate, 328.
Lissorhoptrus simplex, 55–61.
London purple, 194, 195.
Lygæus facetus, 426.
Lygidea mendax, 64.
Lygus invitus, 64.
pratensis, 64, 237.

Macrodactylus subspinosus, 193.
uniformis, 193.
Macrosiphum pisi 457.
Malaria, 31, 32, 33, 34, 36, 93, 100, 104, 217, 370.
Malacosoma americana, 483.
disstria, 483.
Malojillo, 231, 232, 233.
Mamestra picta, 158.
Mangoes, 156.
Maple, 76, 77, 78, 160, 177, 178, 410, 420.
mites, new species, 420–24.
Mare's tail, 238.
Margaropus australis, 223.
Marlatt, C. L., 133–42.
Marlatt scale, 192.
Marmota flaviventer, 209.
May-beetles, 12, 149, 252, 253, 254.
McColloch, J. W., 425–6.
Mealy-bug, 143, 251.
Medical entomology, 27–39.
Mediterranean flour moth, 41.
fruit fly, 134, 135, 140, 149; danger of introducing into United States, 68–73; life history and list of fruits attacked, 399–403; trapping, 347–51.
Megachile addenda, 425.
cleomis, 425.
generosa, 425.
lippiæ, 425.
Megastigmus spermotrophus, 283.
Megilla fuscilabris, 408, 409.
innotata, 149, 257.
Melanconium sacchari, 256.
Melon, 325.
Melon fly, 70.
louse, 190.
Metamasius hemipterus, 251, 252, 253, 256, 450, 454, 455.
sericeus, 248.
Metarrhizium anisopliæ, 247, 452.
Mexican fruit fly, 156, 269.
orange fruit fly, 134.
sugar cane insects, 247–49.
Microbracon juglandis, 144.
vestiticida, 308, 309.
Microsporidium bombycis, 484.
Microterys flavus, 294, 295, 300, 301, 303.
Military grasshopper, 189.
Millett, 362.
Minute Hymenopterous parasites, methods of handling, 341–44.
Miscible oils, 160, 242.
Mocking bird, 365.
Molasses, 157, 236, 401.
Mole-cricket, 12, 252, 255.
Mompha brevivittella, 489.
circumscripta, 489.
pecosella, 489.
Monanthia tabida, 249.
Morelos orange worm, 345.
Morgan, A. C. and Runner, G. A., 226–30.
Morrill, A. W., 185–95; 261–65.

Mosquito, 28, 29, 35, 104, 109, 195, 217, 352, 368.
control in Connecticut, 89–93.
Mountain maple, 422.
Mullen, 239.
Murtfeldt, Mary Esther (Obituary), 288.
Musca domestica, 124.
Mycetobia pallipes, 285.
Myiasis, 373.
Myrmelachista ambigua ramulorum, 150.
Myzus cerasi, 406, 457.
Myzine sexcincta, 255.

Nemorea leucaniæ, 365.
Neosigniphora nigra, 320.
New coccidæ, 142–43.
New destructive insects in New York, 61–65.
New fly trap, 110–12.
Newell, Wilmon, 55–61.
New York apple-tree canker, 178.
Nicotine, 244.
Norway maple, 422, 423, 424.
Notes on inspection, 259–60.
Nysius angustatus, 426.

Oak, 76, 77, 78, 79, 277, 301, 332, 410.
Oats, 117, 278, 362.
Oecanthus nigricornis, 177, 178, 180.
niveus, 177, 178.
quadripunctatus, 177, 178.
Oenothera hewetti, 489.
Oil of citronella, 348.
Oil insecticides on citrus trees and fruits, effects, 161–64.
of pennyroyal, 126.
of tar, 126.
Oligota oviformis, 86.
Onions, 231.
O'Kane, W. C., 153, 159.
Ophthalmia, 31.
Orange, 156, 190, 346, 362, 399.
Orange juice, 401.
Oregon coccinellidæ, 404–7.
Ornithodorus megnini, 223.
Ornix geminatella, 313.
Oryctes rhinoceros, 437.
Oxypleurites dentatus, 422.
dentilobis, 423.

Pamphilius dentatus, 199.
persicum, 199.
Panicum barbinode, 231.
Paracalocoris colon, 64.
Parasitology manuals, 416–18.
Paris green, 236, 440, 478, 479.
Parlatoria blanchardi, 186, 192, 415.
scale, 192.
Parrott, P. J., 61–65.
and Fulton, B. B., 177–80.
Paspalum membranaceum, 57.
Patch, E. M., 316–18.
Peach 156, 157, 177, 178, 180, 190, 193, 237, 238, 240, 263, 345, 400.
ecanium 164.
Peach saw fly, 199.
"Stop Back" and tarnished bug, 237–40.
twig borer, 238.
Peanuts, 362.
Pear, 62, 64, 157, 237, 238, 244, 396.
Psylla, fall spraying, 243–44.
thrips, 62, 65, 66.
Peas, 147, 148.
Pecan, 362.
Pellagra, 34, 113, 366, 374; sand-fly, II, 96–99.
Pepper tree, 191.
Peridroma margaritosa saucia, 192.
Perimeles remus, 249.
Perissopterus javensis, 294, 300, 303.
Peritoma serrulatum, New gall on, 279–80.
Persian walnut, 197, 199.
Peruvian fruit-fly, 345–46.
Petroleum, 126.
fuel oil, 162.
or petroleum products on dormant trees, injuries, 160–61.
Phæniococcus marlattii, 186, 192, 415.
Pheidole megacephala, 330.
Phlebotomus or sand-fly fever, 375.
Philæmatomyia insignis, 475.
Phyllobius argentatus, 63.
calcaratus, 63.
maculicornis, 63.
oblongus, 63.
pyri, 63.
viridicolles, 63.
Phyllocoptes brevisetosus, 423.
constrictus, 423.
magnificus, 423.
minutissimus, 423.
quinquilobus, 424.
splendidus, 424.
trilobis, 424.
variabilis, 424.
Phylloxera vastatrix, 190.
Physokermes insignicola, 298.
Phytalus smithii, 456, 457.
Phytonomus meles, 283.
Picudo of cotton in Peru, 303–12.
Pigeon pea, 325.
Pimpla (Itoplectis) conquisitor, 144–47.
Pine, 76, 78.
Pineapple, 68, 70, 71.
Pinus rigida, 331.
Piojo blanco of cotton, 318–27.
Piroplasma (Babesia), 219.
Pitch pine, 75.
Plague, 32, 35, 378.
Plant lice, 238.
Plasmodium, 219.
falciparum, 367.
vivax, 367.
Plum, 64, 156, 177, 263, 396.
Pocket mice, 224.
Podabrus pruinosus, 406, 457–59.
Pogonomyrmex barbata, 194.

Poison bait, 365.
Poliomyelitis, 30, 32, 34, 35, 103, 104, 105, 106, 107, 108, 109, 110, 113, 366, 375.
Polistes canadensis, 365.
Polydrosus mali, 63.
micans, 63.
sericeus, 63.
Pontia rapæ, 144.
Poplar, 63.
Porthetria dispar, 65.
Potato, 157.
Potato wart, 133.
beetle, 478.
Prenes nero, 257.
Prenolepis fulva, 256.
Prickly pear juice, 401.
Profenusa collaris, 63.
Pronuba yuccasella, 361.
Prospaltella aurantii, 319.
peruviana, 319.
Prune, 396.
Pseudococcus calceolariæ, 445, 450, 454, 455.
sacchari, 12, 251, 252, 253, 255, 455.
Psylla, 66.
Psyllobora tædata, 404.
Pteromalus puparum, 144.
Pulex avium, 84.
Pulvinaria vitis, 295.
Pumpkin, 362.
Pycnoderes quadrimaculatus, 190.

Quail, 365.
Quayle, H. J., 85–88; 290.

Raspberry, 177, 178.
Rats, 378.
Red ant, 330.
bugs, 64.
cedar, 76.
maple, 421, 422, 423, 424.
oak, 77, 177.
spider, 10, 85, 87.
Reduviid bugs, 105.
Remigia repanda, 231, 232, 233, 236, 257.
Rhagoletis cingulata, 79, 80.
fausta, 79, 80.
Rhinoceros beetles, 254, 437–42.
Rhizobius ventralis, 303.
Rhododendron borer, 427.
Rice, 54, 55, 56, 57, 58, 59, 60, 61, 117.
maggots, 59.
water weevil and its control, 55–61.
Riley, W. A., 416–18.
Rind disease, 256.
Ripersia festucæ, 143.
Rocky Mountain spotted fever, 200, 204, 210, 219, 220, 221, 222, 224.
Mountain spotted fever tick, 93–95; 200–11; 218.
Roëntgen rays, effect upon the cigarette beetle, 226–30.
Rose, 191, 192.
Rosenfeld, A. H. and Barber, T. C., 225.
Rosin, 162.
Rosin oil, 162.
Ross, E. H., 429.
Rosy apple aphis, 406, 457.
Rust red flour beetle, 40.
Rutabagas, 362.
Rynchophorus palmarum, 454.

Safro, V. I., 283.
Saissetia oleæ (nigra) 11, 293, 294, 298, 299, 301.
Salix caprea, 63.
Salt, 270.
Salt marsh mosquito, 352, 359.
Sand-fly, 96, 97, 99, 100.
Sanninoidea exitiosa, 193.
San José scale, 9, 66, 191, 263, 266, 267, 404; control, 270–73.
Sap fly, 285.
Sarcophaga georgiana, 365.
Sawtoothed grain beetle, 40.
Scapteriscus didactylus, 12, 252, 255; Cuban parasite of, 330.
Schedius kuvanæ, 298.
Schizoneura lanigera, 190.
Schœne, W. J., 157–58.
Scolia dorsata, 255.
Scolothrips sexmaculatus, 87.
Scrub pine, 331.
Scutellista cyanea, 301.
Scymnus lœwii, 149, 257.
roseicollis, 149, 257.
Severin, H. H. P., 68–73; 399–403.
and Severin, H. C., 347–51.
Shafer, G. D., 160.
Signiphora lutea, 320.
occidentalis, 320.
Silvanus surinamensis, 40.
Simulium vittatum, 97.
Sipha graminis, 149, 257.
Sleeping sickness, 27, 367, 376.
Smaller moth borer, 445, 446, 454.
Smilia misella, 404.
Snout beetles, 54.
Snow-ball aphis, 406.
Snowy tree-crickets, 177.
Soap, 157, 158.
Soft brown scale, 186, 191.
maple, 421.
scale, 293, 294, 300, 301, 302, 303.
Solenopsis geminata, 256, 312, 330.
pylades, 309.
Sorghum, 362, 455.
Southern grass worm, 230, 233, 236, 252, 257.
Soy bean, 362.
Spalaniga muscæ, 124.
Sphæropsis malorum, 178.
Sphenophorus sericeus, 252.
sexguttatus, 253.
Spiders and mites, natural enemies of, 85–88.
Spiny elm caterpillar, 157.

Spirochæta, 219.
Splenetic fever, 32, 33, 34.
Spotted fever, 31, 32, 33, 34, 380.
Squash, 190, 362, 402.
bug, 190.
capsid, 190.
Squirrels, 378.
Stable fly, 30, 100, 105, 112, 113, 115, 116, 217, 459; and infantile paralysis, 101–109; live stock, pest, 112–26; distribution, 459–77.
Stethorus picipes, 87.
Stomoxys calcitrans, 30, 32, 35, 101, 102, 105, 106, 109, 112, 366.
indica, 475.
sugillatrix, 470.
Stop back, 237.
Straw, 125, 126.
String beans, 362, 402.
Sugar beets, 410.
cane, 11, 134, 231, 232, 248, 249, 251, 252, 253, 254, 255, 256, 269, 362, 443–56.
froghopper, 191, 451.
leaf hopper, 455.
mealy bug, 450, 454, 455.
aphid, 149, 257.
beetle, 255.
insects in Porto Rico, 251–57.
mealy-bug, 12, 149, 252, 253, 255, 256.
moth stalk-borer, 12.
scale insect, 257.
Tingid, 247, 249–51.
maple, 285, 421, 422, 423, 424.
solution, 342.
Sulphuric acid, 161, 162.
Sulphur-lime, 9.
Sumac, 178.
Surface, H. A., 265–69.
Swarming of Hemiptera, 426.
Sweet corn, 188.
limes, 156.
potato, 362.
Sycamore, 178.

Tænipoda picticornis, 189.
Tanglefoot, 77.
Tannic acid, 400.
Tar, 440.
Targionia sacchari, 251, 257.
Tarnished plant bug, 237, 238, 239, 240.
Taxodium distichum, 278.
Tenebrioides mauritanicus, 40.
Tepper, J. G. O., 359–61.
Termes flavipes, 284.
morio, 257.
Terminalia catapa, 399.
Tetracha carolina, 364.
Tetranychus bimaculatus, 87.
mytilaspidis, 87.
sexmaculatus, 87.
Tetrastichus asparagi, 144.
Tetrastichus xanthomelænæ, 144.
Tettigonia octolineata, 410.
similis, 257.
Texas fever, 100, 114.
Thermal apparatus, 196–97.
Thriphleps insidiosus, 85.
Thrips, 237.
Ticks, 81, 104, 218, 219, 220, 221, 222, 223, 381.
Tides, rainfall and salt marsh mosquitoes, 352–59.
Timberlake, P. H., 293–303.
Tiphia parellela, 456, 457.
Tobacco, 232.
Tolype velleda, 426.
Tomaspis postica, 247.
varia, 247.
Tomocera californica, 294, 300, 301, 303.
Tomaspis postica, 453.
varia, 444, 451.
Tortrix sacrifaga, 252.
Townsend, C. H. T., 211-25; 244–46; 303–12; 318–27; 345–46.
Tree crickets, 177–80.
Triaspis vestiticida, 308.
Tribolium confusum, 40.
Trichogramma minutum, 341, 447, 454, 456.
pretiosa, 248, 254.
Trifolium, 284.
Trionymus americanus, 143.
californicus, 143.
nanus, 143.
violascens, 143.
Tropical almond, 399, 340, 400.
Trypanosoma, 219.
Trypanosoma cazalboui, 112.
gambiense, 367.
pecaudia, 112.
rhodesiense, 367.
Trypanosomes, 376.
Trypanosomiasis, 27.
Trypeta ludens, 156, 269.
Tsetse flies, 376.
Tuberculosis, 32, 33, 34, 36.
Turnip, 238, 240, 362.
Two elm leaf aphides, 316–18.
Two-spray calendar, success of, 165–73.
Two-stabbed lady-bird beetle, 273.
Typhlocyba comes, 191.
Typhoid fever, 32, 33, 34.
fly, 460.
Typhus fever, 104, 383.

Unspotted tentiform leaf miner of the apple, 313–16.
Urich, F. W., 247–49.

Van Dine, D. L., 251–57.
Variegated cut worm, 192.
Vaseline, 327.
Velvet beans, 362.
Verruga fever, 211–25.
Vetch aphis, 457.

Walnut, 177, 410.
weevil, 197, 198.
weevil, bud moth and a sawfly, 197–99.
Washburn, F. L., 259–60; 327–29.
Wasp sting, effects of, 225.
Watermelon, 362.
Water weevil, 61.
Watson, J. R., 413–14.
Webster, R. L., 180–84; 409–13.
Weevil root-borer, 256, 455.
stalk-borer, 251, 252, 253, 450, 454, 455.
Weiss, H. B., 284, 407–9.
Wells, B. W., 330.
West Indian sugar-cane leaf-hopper, 252, 256.
Western leaf-footed plant bug, 186, 190.
Wheat, 36, 117, 362, 426, 470.
White ants, 284.
fly, 191.
grub, 252.
oak, 77.
pine, 75, 77, 79.
pine blister rust, 133.
scale of cotton, 9.
wash, 273.
Wilson, H. F., 457–59.
Witch hasel, 177.
Wild carrot, 180.
crab, 177.
tobacco, 191.
Willow, 63, 76, 157, 178, 325, 410.
beetle, 158.
Wilt disease, 332.
Woglum, R. S., 290.
Wolcutt, G. N., 443–57.
Woodchuck, 209.
Wooly aphis, 178, 190.

Xerophlæa viridis, 426.
Xyleborus affinis, 248.

Yellow fever, 28, 29, 30, 35, 100, 104, 217, 370.
Yothers, W. W., 161–64.
Yponomeuta malinellus, 64.
padellus, 64.
Yucca aloifolia, flowering and fruit-bearing of, 359–61.

Zebra caterpillar, 158.
Zinc arsenite and arsenate of lead tests, 157–58.
Zizania aquatica, 57.